

DEVELOPMENT AND EVALUATION OF
CENTELLA ASIATICA POWDER BY FREEZE
DRYING: COMPARISON OF EXTRACTION
METHODS ON YIELD AND HYGROSCOPICITY

SITI NUR ARISHA BINTI MUHAMAD
ZULKARNAIN

UNIVERSITI MALAYSIA PAHANG



SUPERVISOR'S DECLARATION

I/We* hereby declare that I/We* have checked this thesis/project* and in my/our* opinion, this thesis/project* is adequate in terms of scope and quality for the award of the degree of Bachelor of Manufacturing Engineering Technology (Pharmaceutical).

(Supervisor's Signature)

Full Name : DR. WAN NURUL HUDA BINTI WAN ZAINAL

Position : SENIOR LECTURER

Date : 10 JANUARY 2019

(Co-supervisor's Signature)

Full Name :

Position :

Date :



STUDENT'S DECLARATION

I hereby declare that the work in this thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Malaysia Pahang or any other institutions.

(Student's Signature)

Full Name : SITI NUR ARISHA BINTI MUHAMAD ZULKARNAIN

ID Number : TD15046

Date : 10 JANUARY 2019

DEVELOPMENT AND EVALUATION OF *CENTELLA ASIATICA* POWDER BY
FREEZE DRYING: COMPARISON OF EXTRACTION METHODS ON YIELD
AND HYGROSCOPICITY

SITI NUR ARISHA BINTI MUHAMAD ZULKARNAIN

Thesis submitted in fulfillment of the requirements
for the award of the degree of
Bachelor of Manufacturing Engineering Technology (Pharmaceutical)

Faculty of Engineering Technology
UNIVERSITI MALAYSIA PAHANG

January 2019

ACKNOWLEDGEMENTS

Primarily, I am grateful to Allah for the good health and wellbeing that were necessary to complete this research.

I would like to express my sincere thanks to Professor Dato' Dr. Engr. Zularisam Bin Ab Wahid, Dean of the Faculty of Engineering Technology, for providing me with all the necessary facilities in completing the research.

I place on record, my sincere thank you to Dr. Wan Nurul Huda Binti Wan Zainal, my supervisor, Dr. Mohd Azmir Bin Arifin and Dr. Raihana Zahirah Binti Edros , senior lecturers in Faculty of Engineering Technology. I am extremely thankful and indebted to them for sharing expertise and valuable guidance and continuous encouragement extended to me.

I am extremely grateful to my parents, Muhamad Zulkarnain Bin Mohd Yunus and Salina Binti Din and my husband, Mustaqim Bin Abd Ghani. Their unconditional love and support has given me strength and courage throughout the journey in completing my degree. I am ever indebted to them for their understanding, patience, and sacrifice, which will always inspire me.

I take this opportunity to express gratitude to all of my friends especially Muhammad Hafiz Bin Yahya, Nur Amalia Binti Ilias, Najihah Binti Salleh, Aimi Zafirah Binti Abd Rahim and Siti Damia Amanda Binti Hamzah that have been sharing so much of their knowledge and opinion to help me.

Last but not least, many thanks to my group members who supported me and to one and all, who directly or indirectly, have lent their hand throughout this venture.

ABSTRAK

Teknik-teknik novel dan pembangunan kaedah terbaharu telah membuka peluang untuk menghasilkan ekstrak sebatian bioaktif semulajadi yang sangat tinggi di samping potensinya untuk merawat dan mencegah pelbagai penyakit. Kajian ini membandingkan peratusan dapatan serbuk *Centella asiatica* yang diperoleh melalui proses pengeringan beku yang diekstrak terlebih dahulu secara akueus melalui kaedah ekstrak konvensional (rebusan dan rendaman) dan kaedah bukan konvensional seperti pengekstrakan bantuan ultrabunyi. Kestabilan fizikal serbuk kemudiannya diuji melalui ujian higroskopik. Ujian ini menggunakan teknik analisis serapan gravimetrik dengan sedikit modifikasi. Setiap sampel serbuk melalui pra-rawatan pada 45 °C, 10% RH diikuti dengan proses ekuilibrasi pada 25 °C, 80% RH dan seterusnya dikategorikan mengikut European Pharmacopeia. Pengekstrakan bantuan ultrabunyi menghasilkan dapatan serbuk terbanyak (11.65%) diikuti oleh rendaman (10.48%) dan rebusan (10.03%). Setelah 24 jam dianalisa, ketiga-tiga sampel diklasifikasikan sebagai sangat higroskopik dengan peratusan serapan air melebihi 15%. Peratusan serapan air tertinggi adalah melalui rebusan (71.31%), diikuti oleh rendaman (71.28%) dan pengekstrakan bantuan ultrabunyi (68.71%).

ABSTRACT

Novel techniques and the development of new methods has provided an opportunity to obtain highly purified natural bioactive compound extracts with potential for the treatment and prevention of various diseases. This research compares the percentage yield of *Centella asiatica* obtained through freeze drying that was aqueous-extracted earlier by conventional extraction method (decoction and infusion) and an advanced non-conventional extraction method; ultrasound-assisted extraction (UAE). The physical stability of the powder obtained was then tested by hygroscopicity test. The studies were conducted using gravimetric sorption analysis method with slight modification. Each of the powder samples was subjected to the pretreatment at 45 °C, 10% RH followed by equilibrating at 25 °C, 80% RH and individually categorized as per European Pharmacopeia. UAE generated the highest powder yield (11.65%) followed by infusion (10.48%) and decoction (10.03%). After 24 hours of hygroscopicity analysis, all three samples were found to be very hygroscopic since their respective percentage of water sorption was more than 15%. The highest percentage of water sorption was by decoction (71.31%), followed by infusion (71.28%) and lastly UAE (68.71%).

TABLE OF CONTENT

DECLARATION

TITLE PAGE

ACKNOWLEDGEMENTS ii

ABSTRAK iii

ABSTRACT iv

TABLE OF CONTENT v

LIST OF TABLES viii

LIST OF FIGURES ix

LIST OF SYMBOLS xi

LIST OF ABBREVIATIONS xii

CHAPTER 1 INTRODUCTION 1

1.1 Background Study 1

1.2 Problem Statement 3

1.3 Research Objective 4

1.4 Scope of Study 4

CHAPTER 2 LITERATURE REVIEW 5

2.1 Introduction to *Centella Asiatica* 5

2.1.1 Botanical Aspects and Macroscopic of *C. asiatica*, *H. bonariensis* and *H. sibthorpioides* 6

2.1.2 Microscopic Evaluation of *C. asiatica* Leaves 12

2.1.3 Microscopic Evaluation of *C. asiatica* Petiole 14

2.2	Extraction	16
2.3	Extraction Solvent	16
2.4	Extraction Methods	19
2.4.1	Decoction	19
2.4.2	Infusion	20
2.4.3	Ultrasound-assisted Extraction (UAE)	21
2.5	Freeze Drying	22
2.5.1	History and Development	22
2.5.2	Characteristics of Freeze Drying	22
2.5.3	Eutectic/Collapse Temperature	24
2.5.4	Primary Drying	25
2.5.5	Secondary Drying	25
2.6	Hygroscopicity	26
CHAPTER 3 METHODOLOGY		28
3.1	Material/Chemical	29
3.2	Equipment	29
3.3	Moisture Content of Fresh <i>C. asiatica</i>	30
3.4	Preparation of <i>C. asiatica</i> Aqueous Extract	30
3.4.1	Plant Materials	30
3.4.2	Decoction	30
3.4.3	Infusion	31
3.4.4	Ultrasound- assisted Extraction (UAE)	32
3.5	Basic Operation of Freeze Dryer	33
3.6	Freezing of <i>C. asiatica</i> Extract	35
3.7	Primary Drying of <i>C. asiatica</i> Extract	35

3.8	Secondary Drying of <i>C. asiatica</i> Extract	35
3.9	Yield of <i>C. asiatica</i> Powder	36
3.10	Determination of Extractable Solids	36
3.11	Hygroscopicity Study	36
CHAPTER 4 RESULTS AND DISCUSSION		38
4.1	Moisture content of fresh <i>C. asiatica</i> plant	38
4.2	Yield of <i>C. asiatica</i> powder	39
4.3	Extractable solids	42
4.4	Hygroscopicity study	42
CHAPTER 5 CONCLUSION		49
5.1	Conclusion	49
5.2	Recommendation	50
REFERENCES		50
APPENDIX A		59
APPENDIX B		61

LIST OF TABLES

Table 2.1	Antioxidant phytoconstituents of 100% ethanol extract of <i>C.asiatica</i> (100% E Ex) and water extract of <i>C. asiatica</i> (H ₂ O Ex)	17
Table 2.2	TPC and TAC of herbal plant (<i>Tilia cordata</i>) extracts obtained by infusion and UAE	20
Table 2.3	Hygroscopicity categorization as per Callahan et al. and Ph. Eur.	27
Table 3.1	Material list and quantity	29
Table 3.2	Equipment list	29
Table 4.1	Percentage of extractable solids obtained from 10 ml extracts from each extraction methods presented as % (g/ 10 ml)	42
Table 4.2	Mass losses during pre-treatment of powders obtained from different extraction methods presented in % (<i>m/m</i>)	44
Table 4.3	Mass gained during gravimetric sorption analysis method of powder samples obtained from different extraction methods presented in % (<i>m/m</i>)	46
Table 4.4	<i>C. asiatica</i> powder categorization per Ph. Eur.	47

LIST OF FIGURES

Figure 1.1	<i>Centella Asiatica</i>	1
Figure 2.1	(a) <i>C. asiatica</i> , (b) Parts of <i>C. asiatica</i> , (c) <i>C. asiatica</i> : Leaf (stereo microscopic view x 1), (d) <i>C. asiatica</i> : Stem (Stereo microscopic view x 1), (e) <i>C. asiatica</i> : Inflorescence (stereo microscopic view x 1), (f) <i>C. asiatica</i> : Inflorescence (stereo microscopic view x 1) and (g) <i>C. asiatica</i> : Inflorescence (stereo microscopic view x 2)	7
Figure 2.2	(a) <i>H. bonariensis</i> clone, showing individual ramets and lateral runner growth pattern. Youngest ramets are produced at the tip (indicated by arrow), (b) Individual <i>H. bonariensis</i> ramets showing leaves of typical orientations (leaf angles and azimuths), (c) Single <i>H. bonariensis</i> runner (1) with individual ramet (2) consisting of a single broad leaf and individual root system	9
Figure 2.3	(a) <i>H. sibthorpioides</i> : Leaf shape, (b) Single <i>H. sibthorpioides</i> runner consisting of a single broad leaf, (c) Single <i>H. sibthorpioides</i> runner with individual root system, (d) <i>H. sibthorpioides</i> : Inflorescence (stereo microscopic view x 1) and (e) <i>H. sibthorpioides</i> : Inflorescence (stereo microscopic view x 2)	10
Figure 2.4	(a) <i>C. asiatica</i> : T.S. of lamina through midrib (x 4), (b) <i>C. asiatica</i> : T.S. of lamina through midrib (x 10), (c) <i>C. asiatica</i> : A portion of midrib is enlarged (x 40) and (d) <i>C. asiatica</i> : T.S. of lamina- a portion enlarged (x 40)	13
Figure 2.5	(a) <i>C. asiatica</i> : T.S. of petiole (x 2), (b) <i>C. asiatica</i> : T.S. of petiole (x 4), and (c) <i>C. asiatica</i> : a portion of petiole enlarged (x10)	15
Figure 2.6	Mechanism of UAE method	21
Figure 2.7	Phase diagram of water	23
Figure 2.8	Freeze drying steps	23
Figure 3.1	Flowchart of the entire process	28
Figure 3.2	Decoction of <i>C. asiatica</i>	31
Figure 3.3	Infusion of <i>C. asiatica</i>	32
Figure 3.4	Ultrasound-assisted extraction of <i>C. asiatica</i>	33
Figure 3.5	Freeze drying process via VirTis BenchTop Pro with Omnitronics-8L	34

Figure 4.1	Curve of percentage moisture content loss on fresh <i>C. asiatica</i> during drying treatment in 30 hours	39
Figure 4.2	Crude powders obtained post-freeze dry of 46.3 g dried <i>C. asiatica</i> in each extraction methods presented in % yield (g/46.3 g)	40
Figure 4.3	Structural classification of water in solids	44

LIST OF SYMBOLS

cm	Centimetre
°C	Degree Celsius
G	Gram
hr	Hour
kcal	Kilo Calories
kg	Kilogram
kHz	Kilo Hertz
MHz	Mega Hertz
<i>m/m</i>	Mass per Mass
µg	Microgram
µm	Micrometre
mg	Milligram
ml	Millilitre
mm	Millimetre
mm Hg	Millimetre Mercury
mTorr	MilliTorr
mol	Mole
%	Percentage
w/v	Weight per Volume
w/w	Weight per Weight

LIST OF ABBREVIATIONS

AE	Ascorbic acid Equivalent
T _c	Collapse Temperature
E Ex	Ethanol Extract
Ph. Eur.	European Pharmacopeia
T _e	Eutectic Temperature
ES	Extractable Solids
GAE	Gallic Acid Equivalent
GAG	Glycosaminoglycan
M _{GSA}	Mass Gravimetric Sorption Analysis
M _{PT}	Mass Pre-treatment
PE	Pyrogallol Equivalent
QE	Quercetin Equivalent
RH	Relative Humidity
TE	Tannic acid Equivalent
TAC	Total Antioxidant Capacities
TFC	Total Flavonoid Content
TPC	Total Phenolic Content
TE	Trolox Equivalent
UAE	Ultrasound-Assisted Extraction
VH	Very Hygroscopic
H ₂ O Ex	Water Extract
XOI	Xanthine Oxidase Inhibitory

CHAPTER 1

INTRODUCTION

1.1 Background Study

Centella asiatica as shown in Figure 1.1, commonly known as centella, Asiatic pennywort or Gotu kola, is a herbaceous, frost-tender perennial plant belonging to the family Apiaceae. Easily grow and found in the most tropical and subtropical countries, *C. asiatica* is used for medicinal purposes (Singh and Singh, 2002). *C. asiatica* or Gotu kola should not be confused with kola nut as it does not contain any caffeine and has not been shown to have stimulant properties (Niamnuy et al., 2013). There are several important bioactive compounds present in *C. asiatica* such as triterpene saponins, phenolic compounds, vitamins, minerals, free amino acids, and polyacetylenic compounds (Niamnuy et al., 2013). All of these compounds are the major ingredients that are responsible for *C. asiatica* therapeutic effects.



Figure 1.1 *Centella Asiatica* (Source: Sudhakaran, 2017)

In the nineteenth century, *C. asiatica* and its extracts were incorporated into the Indian pharmacopoeia, wherein in addition to wound healing, it was recommended for various skin conditions treatment (Gohil, Patel and Gajjar, 2010; Gohil, Patel and Gajjar, 2012). *C. asiatica* extracts contain 70% triterpenic acids mainly two triterpene glycosides (asiaticoside and madecassoside) and corresponding aglycones (asiatic acid and madecassic acid) which exerts normalizing action on metabolism of connective tissues. It also enhances tissue integrity by stimulating glycosaminoglycan (GAG) synthesis without promoting excessive collagen synthesis or cell growth (Pizzorno, Murray and Joiner-Bey, 2016; Govarathanan et al., 2015). However, it is inconvenience for some users to consume it as it is. Thus, the *C. asiatica* plant is extracted before it is powdered and encapsulated to enhance its bioavailability, stability and shelf life (Rivas et al., 2017).

Extraction of botanicals and herbal sample is the crucial first step prior to analysis of bioactive compounds present in the preparations (Sasidharan et al., 2011). Various solvents can be used to extract the *C. asiatica* which includes methanol (Govarathanan et al., 2015), ethanol (Dewi and Maryani, 2015) and distilled water (Cheng et al., 2004). Different solvents will give different quantity of yield obtained. Due to unmatched availability in chemical diversity, pure compounds or standardized plant extracts provides unlimited opportunities for new drug discoveries (Cos et al., 2006). The extraction basic steps are pre-washing the sample, drying or freeze drying the plant material, grinding to obtain a homogenous sample and enhancing the kinetics of analytic extraction and also increasing the contact of sample surface with the solvent system. During the extract preparation, proper actions must be taken to assure that there is no loss, distortion or denaturation of potential active constituents of the plant samples (Sasidharan et al., 2011).

Drying is a common method of preservation, applied by removing the water from the plant. Convection drying and freeze drying are the two most popular drying methods. Conventional drying such as oven drying is a user-friendly and very low cost method compared to freeze drying, which is a lengthy procedure and requires higher cost (Vashisth, Singh and Pegg, 2011). Previous research showed that there are major quality (nutrient compound and colour) differences of the final product after conventional drying compared to freeze dried products (Wojdylo et al., 2014).

Freeze drying is conducted to remove the solvent in such a way that the sensitive molecular structure of the active substance in the plant is least disturbed (Alexeenko, Ganguly and Nail, 2009). It is more suitable for products sensitive to temperature or to high residual moisture content (Touzet et al., 2018).

1.2 Problem Statement

The raw, unprocessed *C. asiatica* plant have a shorter shelf life and physically unstable. The leaves are easily wilted upon exposure to sunlight and its bioactive components degraded due to oxidation which leads to physical changes of the plant. *C. asiatica* fresh plant is very likely susceptible towards microbial growth due to high moisture content especially in leaves and stems (Brinkhaus et al., 2000). Besides that, it may be inconvenient for patient to consume it raw since some of them may not like to eat vegetables or herbs. This study is conducted to overcome this problem by converting the *C. asiatica* into powders. A study shows that *C. asiatica* lost a high amount of phenolic compounds (total phenolic content (TPC) and total flavonoid content (TFC) when extracted with ethanol of concentration 80% and above (Chew et al., 2011). In addition, the safety consumption of oral drug with active ingredients extracted using alcohol solvents is probably the main issue to deal with (Man and Choo, 2018). In other aspect, conventional extraction methods (maceration, decoction, infusion etc.) are the most convenient and inexpensive as it does not require any advanced equipment (Easmin et al., 2014). However, commercial extraction methods such as microwave-assisted extraction, ultrasound-assisted extraction and super critical fluid extraction have been proposed to extract bioactive compounds from plants with higher recovery capacity (Caldas et al., 2018). Hence, the solvent as well as the extraction methods used must be optimized to obtain the highest yield of *C. asiatica* powder that is physically more stable.

REFERENCES

- Alexeenko, A.A., Ganguly, A., and Nail, S.L. (2009). Computational analysis of fluid dynamics in pharmaceutical freeze-drying. *Journal of Pharmaceutical Sciences*, Vol. 98. Issue 9. 3483-3494
- Allada, R., Maruthapillai, A., Palanisamy, K., and Chappa, P. (2016). Hygroscopicity categorization of pharmaceutical solids by gravimetric sorption analysis: A systematic approach. *Asian Journal of Pharmaceutics*, Vol. 10. Issue 4. 279-286
- Aramouni, F., and Deschenes, K. (2015). Methods for developing new food products. DEStech Publications Inc. 154-156
- Atha, D. (2017). Hydrocotyle Sibthorpioides and H. Batrachium (Araliaceae) new for New York State. *Phytoneuro*, Vol. 56. 1–6
- Belachew, T., and Lindstrom, D. (2015). Practical techniques of dehydration of foods. Koros Press Editorial Board. 85-88
- Brinkhaus, B., Lindner, M., Schuppan, D., and Hahn, E. G. (2000). Chemical, pharmacological and clinical profile of the East Asian medical plant Centella Asiatica. *Phytomedicine*, Vol. 7(5). 427-448
- Caldas, T. W., Mazza, K. E. L., Teles, A. S. C., Mattos, G. N., Brigida, A. I. S., Conte-Junior, C. A., Borguini, R. G., Godoy, R. L. O., Cabral, L. M. C., Tonon, R. V. (2018). Phenolic compounds recovery from grape skin using conventional and nonconventional extraction methods. *Industrial Crops & Products*, Vol. 111. 86-91
- Callahan, J.C., Cleary, G.W., Elefant, M., Kaplan, G., Kensler, T., and Nash, R.A. (1982). Equilibrium moisture content of pharmaceutical excipients. *Drug Development and Industrial Pharmacy*, Vol. 8. Issue 3. 355-369
- Castro-López, C., Ventura-Sobrevilla, J.M., González-Hernández, M.D., Rojas, R., Ascacio-Valdés, J.A., Aguilar, C.N., and Martínez-Ávila, G.C.G. (2017). Impact of extraction techniques on antioxidant capacities and phytochemical composition of polyphenol-rich extracts. *Food Chemistry*, Vol. 237. 1139-1148

- Chemat, F., Rombaut, N., Sicaire, A-G., Meullemiestre, A., Fabiano-Tixier, A-S. , and AbertVian, M. (2016). Ultrasound assisted extraction of food and natural products. mechanisms, techniques, combinations, protocols and applications A Review. *Ultrasonics Sonochemistry*
- Chen, D. (2009). Hygroscopicity of pharmaceutical crystals. Faculty of Graduate School of the University of Minnesota
- Cheng, C. L., Guo, J. S., Luk, J., and Leung Koo, M. W. (2004). The healing effects of Centella extract and asiaticoside on acetic acid induced gastric ulcers in rats. *Life Sciences*, Vol. 74. Issue 18. 2237-2249
- Chew, K. K., Ng, S. Y., Thoo, Y. Y., Khoo, M. Z., Wan Aida, W. M., and Ho, C. W. (2011). Effect of ethanol concentration, extraction time and extraction temperature on the recovery of phenolic compounds and antioxidant capacity of Centella Asiatica extracts. *International Food Research Journal*, Vol. 18. 571-578
- Cittan, M., Altuntas, E., and Celik, A. (2018). Evaluation of antioxidant capacities and phenolic profiles in Tilia cordata fruit extracts: A comparative study to determine the efficiency of traditional hot water infusion method. *Industrial Crops & Products*, Vol. 122, 553-558
- Cong-Cong, X., Bing, W., Yi-Qiong, P., Jian-Sheng, T, and Tong, Z. (2017). Advances in extraction and analysis of phenolic compounds from plant materials. *Chinese Journal of Natural Medicines*, Vol. 15 (10). 721-731
- Cos, P., Vlietinck, A.J., Berghe, D. V., and Maes, L. (2006). Anti-infective potential of natural products: how to develop a stronger in vitro ‘proof-of-concept’. *Journal of Ethnopharmacology*, Vol. 106. Issue 3. 290-302
- De Bruijn, J., Rivas, F., Rodriguez, Y., Loyola, C., Flores, A., Melin, P., and Borquez, R. (2016). Effect of vacuum microwave drying on the quality and storage stability of strawberries. *J. Food Process. Preservation*, Vol. 40. 1104-1115
- Dent, M., Dragovic-Uzelac, V., Penic, M., Brncic, M., Bosiljkov, T and Levay, B. (2013). The effect of extraction solvents, temperature and time on the composition and mass fraction of polyphenols in dalmatian wild sage (*Salvia officinalis* L.) extracts. *Food Technology and Biotechnology*, Vol. 51. No 1. 84–91

- Dewi, R.T., and Maryani, F. (2015). Antioxidant and α -glucosidase inhibitory compounds of *Centella Asiatica*. *Procedia Chemistry*, Vol. 17. 147-152
- Easmin, MT. S., Sarker, M. Z. I., Ferdosh, S., Shamsudin, S. H., Yunus, K., Uddin, M. S., Sarker, M. M. R., Akanda, M. J. H., Hossain, M. S., and Khalil, HSP. A. (2014). Bioactive compounds and advanced processing technology: *Phaleria macrocarpa* (sheff.) Boerl, A review. *Journal of Chemical Technology & Biotechnology*
- Franks, F.(2008). Freeze-drying of pharmaceuticals and biopharmaceuticals: principles and practice. Royal Society of Chemistry Publishing. 105-128
- Gohil K.J., Patel J.A., Gajjar A.K. (2012). A pharmacological investigation of the potential herbal drug interaction between *Centella Asiatica* and imipramine in rats and mice. *Asian Pacific Journal of Tropical Biomedicine*. 1-11
- Gohil K.J., Patel J.A., Gajjar A.K.. (2010). Pharmacological review on *Centella asiatica*: A potential herbal cure-all. *Indian Journal of Pharmaceutical Sciences*, Vol. 72. No 5. 546-556
- Govarthanan, M., Rajinikanth, R., Kamala-Kannan, S., and Selvankumar, T. (2015). A comparative study on bioactive constituents between wild and in vitro propagated *Centella Asiatica*. *Journal of Genetic Engineering and Biotechnology*, Vol. 13. Issue 1. 25-29
- Joesting, H. M., Sprague, M. O., and Smith W. K. (2012). Seasonal and diurnal leaf orientation, bifacial sunlight incident, and leaf structure in the sand dune herb *Hydrocotyle Bonariensis*. *Environmental and Experimental Botany*, Vol. 75. 195-203
- Karathanos, V. T., Anglea, S. A., and Karel, M. (1996). Structural collapse of plant materials during freeze-drying. *Journal of Thermal Analysis*, Vol. 47. Issue 5. 1451–1461
- Katalinic, V., Milos, M., Kulisic, T., and Jukic, M. (2006). Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food Chemistry*, Vol. 94. 550-557
- Khatib, A., H, M., Shaari, K., Abas, F., Shitan, M., Kneer, R., Neto, V., and Lajis, H.N.(2012). Discrimination of three pegaga (*Centella*) varieties and determination of growth-lighting effects on metabolites content based on the chemometry of 1h nuclear magnetic resonance spectroscopy. *Journal of Agricultural and Food Chemistry*, Vol. 60. No 1 . 410-417

- Labconco. (2004). A guide to freeze drying for the laboratory. Retrieved from <http://www.thermofishersci.in/lit/Labconco%20-%20A%20laboratory%20guide%20to%20freeze%20drying.pdf>
- Lemus-Mondaca, R., Vega-Galvez, A., Rojas, P., Stucken, K., Delporte, C., Valenzuela-Barra, G., Jagus, R. J., and Agüero, M. V. (2018). Antioxidant, antimicrobial and anti-inflammatory potential of *Stevia rebaudiana* leaves: effect of different drying methods. *Journal of Applied Research on Medicinal and Aromatic Plants*
- Levi, G., and Karel, M. (1995). Volumetric shrinkage (collapse) in freeze-dried carbohydrates above their glass transition temperature. *Food Research International*, Vol. 28. Issue 2. 145-151
- Llyod, P.J., and Wyk, J.V. (2012). Enhancing extraction processes in the food industry. CRC Press. 1-22
- LucidCentral. (2018). Herbal plant. Retrieved from https://keys.lucidcentral.org/keys/v3/bonsai/en/key/weeds_in_bonsai_en/Media/Html/desc_Hydrocotyle_sibthorpioides_var_sibthorpioides.html
- Ma, Q., Santhanam, R. K., Xue, Z., Guo, Q., Gao, X., and Chen, H. (2018). Effect of different drying methods on the physicochemical properties and antioxidant activities of mulberry leaves polysaccharides. *International Journal of Biological Macromolecules*
- Malaysia Meteorological Department. (n.d). Maklumat ramalan cuaca. Retrieved from <http://www.met.gov.my/home>
- Man, F., and Choo, C.Y. (2018). Safety assessment of standardized aqueous *Brucea Javanica* extract in rats. *Journal of Ethnopharmacology*, Vol. 215. 21-26
- Maulidiani, M., Abas, F., Khatib, A., Shaari, K., and Lajis, N.H. (2014). Chemical characterization and antioxidant activity of three medicinal apiaceae species. *Industrial Crops and Products*, Vol. 55. 238–247
- Newman, A.W., Reutzel-Edens, S.M. and Zografi, G. (2008). Characterization of the “hygroscopic” properties of active pharmaceutical ingredients. *Journal of Pharmaceutical Sciences*, Vol. 97. No 3. 1047-1059

- Ngo, T. V., Scarlett, C. J., Bowyer, M. C., Ngo, P. D., and Vuong, Q. V. (2017). Impact of different extraction solvents on bioactive compounds and antioxidant capacity from the root of *Salacia chinensis* L. *Journal of Food Quality* 2017
- Niamnuy, C., Charoenchaitrakool, M., Mayachiew, P., and Devahastin, S. (2013). Bioactive compounds and bioactivities of *Centella asiatica* (L.) Urban prepared by different drying methods and conditions. *International Journal of Drying Technology*, Vol. 31. Issue 16. 2007-2015
- Oreopoulou, A., Tsimogiannis, D., and Oreopoulou, V. (n.d). Extraction of polyphenols from aromatic and medicinal plants: An overview of the methods and the effect of extraction parameters. *National Technical University of Athens*
- Panja, P. (2017). Green extraction methods of food polyphenols from vegetable materials. *Food Science*, Vol. 10
- Pferschy-Wenzig, E. M., and Bauer, R. (2015). The relevance of pharmacognosy in pharmacological research on herbal medicinal products. *Epilepsy & Behavior*, Vol. 52. 344-362
- Pizzorno, J. E., Murray, M. T., and Joiner-Bey, H. (2016). The clinician's handbook of natural medicine. Churchill Livingstone. Edition 3. 960-966
- PlantNET. (1992). *Hydrocotyle bonariensis* Lam. Retrieved from <http://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=sp&name=Hydrocotyle~bonariensis>
- Plengmuankhae, W., and Tantitadapitak, C. (2015). Low temperature and water dehydration increase the levels of asiaticoside and madecassoside in *Centella Asiatica* (L.) Urban. *South African Journal of Botany*, Vol. 97. 196-203
- Prosapio, V., Norton, I., and De Marco, I. (2017). Optimization of freeze-drying using a life cycle assessment approach: strawberries' case study. *Journal of Cleaner Production*, Vol. 168. 1171-1179
- Pyper, J. W. (1985). The determination of moisture in solids: A selected review. *Analytica Chimica Acta*, Vol. 170. 159-175

- Rahman, M., Hossain, S., Rahaman, A., Fatima, N., Nahar, T., Uddin, B., and Ahmed Basunia, M. (2013). Antioxidant activity of *Centella asiatica* (Linn.) Urban: impact of extraction solvent polarity. *Journal of Pharmacognosy and Phytochemistry*, Vol. 1. Issue 6
- Ratti, C. (2001). Hot air and freeze-drying of high-value foods: a review. *Journal of Food Engineering*, Vol. 49. Issue 4. 311-319
- Rivas, C. J. M., Tarhini, M., Badri, W., Miladi, K., Greige-Gerges, H., Nazari, Q. A., Rodríguez, S. A. G., Román, R. A., Fessi, H., and Elaissari, A. (2017). Nanoprecipitation process: from encapsulation to drug delivery. *International Journal of Pharmaceutics*, Vol. 532. Issue 1. 66-81
- Ruenroengklin, N., Zhong, J., Duan, X., Yang, B., Li, J., and Jiang, Y. (2008). Effects of various temperatures and pH values on the extraction yield of phenolics from litchi fruit pericarp tissue and the antioxidant activity of the extracted anthocyanins. *International Journal of Molecular Sciences*, Vol. 9. No 7. 1333-1341
- Sabaragamuwa, R., Perera, C. O., and Fedrizzi, B. (2018). *Centella Asiatica* (Gotu Kola) as a neuroprotectant and its potential role in healthy ageing. *Trends in Food Science & Technology*, Vol. 79. 88-97
- SADC: South African Development Community. (2004). SADC guideline for stability testing. Retrieved from http://www.ich.org/fileadmin/Public_Web_Site/ABOUT_ICH/Organisation/SADC/Guideline_for_Stability_Studies.pdf
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K. M., and Yoga Latha, L. (2011). Extraction, isolation and characterization of bioactive compounds from plants extracts. *Afr J Tradit Complement Altern Med*, Vol. 8. No 1. 1–10
- SEPPIC. (n.d). Challenges and solutions for moisture sensitive API formulation. Retrieved from <https://www.seppic.com/pharmaceuticals/ingredients>
- Singh, P., and Singh, J. S. (2002). Recruitment and competitive interaction between ramets and seedlings in a perennial medicinal herb, *Centella Asiatica*. *Basic and Applied Ecology*, Vol. 3. Issue 1. 65-76

- Stanisavljevic, D.M., Dordevic, S.M., Ristic, M.S., Velickovic, D.T., and Randelovic, N.V. (2010). Effects of different drying methods on the yield and the composition of essential oil from herb *Mentha Longifolia* (L.) Hudso. *Biologica Nyssana*, Vol. 1. No 1-2. 89-93
- Sudhakaran, M. V. (2017). Botanical pharmacognosy of *Centella asiatica* (Linn.)Urban. *Pharmacognosy Journal*, Vol. 9(4). 546-558
- Tan, S.H. (2010). Production of flavonoids in pegaga (*Centella Asiatica* L. Urban) cell suspension culture. PhD thesis, Universiti Putra Malaysia.
- Touzet, A., Pfefferléa, F., Wel, P.V.D., Lamprecht, A., and Pellequer, Y. (2018). Active freeze drying for production of nanocrystal-based powder: A pilot study. *International Journal of Pharmaceutics*, Vol. 536. Issue 1. 222-230
- USDA-NRCS: United States Department of agriculture-natural resources convention service. (n.d) *Hydrocotyle Bonariensis* Comm. ex Lam. Retrieved from <https://plants.usda.gov/core/profile?symbol=hybo>
- Vashisth, T., Singh, R.K., and Pegg, R.B. (2011). Effects of drying on the phenolics content and antioxidant activity of Muscadine Pomace. *LWT - Food Science and Technology*, Vol. 44. Issue 7. 1649-1657
- VirTis SP Scientific. (2014). BenchTop pro with Omnitronics-8L: freeze dryer operator's manual
- Vongsak, B., Sithisarn, P., Mangmool, S., Thongpraditchote, S., Wongkrajang, Y., and Gritsanapan, W. (2013). Maximizing total phenolics, total flavonoids contents and antioxidant activity of *Moringa Oleifera* leaf extract by the appropriate extraction method. *Industrial Crops and Products*, Vol. 44. 566-571.
- Wang, L. and Weller, C.L. (2006). Recent advances in extraction of nutraceuticals from plants. *Trends In Food Science & Technology*, Vol. 17. 300-312
- Wojdylo, A., Figiel, A., Lech, K., Nowicka, P., and Oszmiański, J. (2014). Effect of convective and vacuum–microwave drying on the bioactive compounds, color, and antioxidant capacity of sour cherries. *Food and Bioprocess Technology*, Vol. 7. Issue 3. 829-841

- Yahya, N.A., Attan, N. and Abdul Wahab, R. (2018). An overview of cosmeceutically relevant plant extracts and strategies for extraction of plant-based bioactive compounds. *Food and Bioproducts Processing*, Vol. 112. 69-85
- Zainol, M.M.K., Abdul-Hamid, A., Abu Bakar, F., and Pak Dek, S. (2009). Effect of different drying methods on the degradation of selected flavonoids in *Centella Asiatica*. *International Food Research Journal*, Vol. 16. 531-537
- Zainol, N.A., Voo, S.C., Sarmidi, M.R., and Aziz, R.A. (2008). Profiling of *Centella Asiatica* (L.) Urban Extract . *The Malaysian Journal of Analytical Sciences*, Vol. 12. No 2. 322 -327